

Dosage Charts



Acetaminophen (Tempra, Panadol, Generic)

Dosage: 15mg/kg every 4-6 hours as needed

	7-11 lbs.	12-17 lbs.	18-23 lbs.	24-35 lbs.	36-47 lbs.	48-59 lbs.	60-70 lbs.	71-100 lbs.
Syrup Elixir (Infants Tylenol or Children's Tylenol® Oral Suspension) 160 mg/5ml	1.25 ml	2.5 ml	3.75 ml	5 ml	7.5 ml	10 ml	12.5 ml	15 ml
Children's Chewable 160 mg tabs				1 tab	1½ tab	2 tabs	2½ tabs	3 tabs
Suppository 80 mg		1 Supp.	1½ Supp.	2 Supp.				
Suppository 120 mg			1 Supp.	1½ Supp.				
Suppository 325 mg				½ Supp.	¾ Supp.	1 Supp.		

	47-69 lbs.	70-95 lbs.	95+ lbs.
Adult 325 mg	1 tab	1½ tabs	2 tabs
Adult 500 mg		1 tab	1 tab

IBUPROFEN (Motrin®, Advil®) * Not recommended for under six months

Dosage: 10mg/kg every 6-8 hours as needed

	12-17 lbs.	18-23 lbs.	24-35 lbs.	36-47 lbs.	48-59 lbs.	60-71 lbs.	72-95 lbs.
Infant Drops 50 mg/1.25 ml	1.25 ml	1.875 ml	2.5 ml	3.75 ml	5 ml		
Suspension 100 mg/5 ml	2.5 ml	3.75 ml	5 ml	7.5 ml	10 ml	12.5 ml	15 ml
Chewable 50 mg			2-3 tabs	3-4 tabs	4-5 tabs	5-6 tabs	6-7 tabs
Chewable 100 mg			1-1½ tabs	1½-2 tabs	2-2½ tabs	2½-3 tabs	3-3½ tabs
Caplets 100 mg			1-1½ tabs	1½-2 tabs	2-2½ tabs	2½-3 tabs	3-3½ tabs
Adult Tabs 200 mg					1 tab	1-1½ tabs	1½-2 tabs

Remember to check active ingredients in cough/cold medicines, as they may contain Acetaminophen or Ibuprofen and should not be given with the above medication.

Always double check the concentration of your medicine bottle with our guidelines for dosing as some generic forms may be different. Always use the dropper that came with the bottle for the most accurate dosing. Call if you have questions.